Pneumoconiosis and Silicosis: Recent Trend and Public Health Responses

Zhongqi Chen*

Department of Pharmacology and Toxicology, University of Toronto, Toronto, M5S1A1, Canada

*Corresponding author: zhongqi.chen@mail.utoronto.ca

Abstract. Pneumoconiosis, a collective term for lung diseases resulting from the inhalation of specific dusts, presents significant global public health and economic challenges1. This is particularly prevalent in traditional sectors like mining and metal industries, and emerging industries such as nuclear fuel processing. Notably, nations like China and India have over 20 million workers at risk. While awareness around pneumoconiosis has been prevalent for decades, its status as a leading occupational disease, especially in regions like China, indicates a deficiency in current mitigation policies. This essay delves into the present diagnostic and treatment measures for pneumoconiosis, its prevention strategies, worldwide trends, and the existing policies from governmental and private entities, highlighting areas for enhancement. The global resurgence of pneumoconiosis underscores the pressing need for a thorough reassessment of workplace regulations and the creation of novel standards tailored to emergent industries. Comprehensive research aimed at identifying air pollutants and associated risks in new industry workspaces is imperative. Collaborative endeavors from governmental and private sectors should focus on enhanced protective gear, workplace safety education, and comprehensive medical insurance. Innovative strategies, like forecasting potential workplace pollutants, can bridge the gap between policy enforcement and safety measures, aiming to reduce the influence of this incapacitating ailment.

Keywords: Pneumoconiosis; policy; public health.

1. Introduction

Pneumoconiosis exerts a profound influence on global public health and economic landscapes, especially within long-established sectors such as the mining and metallurgical industries, and in burgeoning domains like nuclear fuel production and specialized equipment fabrication. The designation "pneumoconiosis" refers to an array of pulmonary diseases instigated by the inhalation of specific particulate matter [1-4]. To illustrate, silicosis, a subtype of pneumoconiosis culminating in pulmonary fibrosis, arises from the inhalation of crystalline silica particulates, a common occurrence in industries such as glass production and silica extraction [5]. For numerous decades, pneumoconiosis has stood as a salient public health conundrum. In nations like China and India alone, in excess of 20 million workers confront the perils associated with such diseases [4]. A sizable fraction of this workforce is entrenched in coal mining and thermoelectric power generation sectors [3,4]. Compounding this issue is the escalating count of individuals in nascent industries, including nuclear fuel refinement and specialized apparatus fabrication, who are afflicted by pneumoconiosis [3]. Notwithstanding its sustained visibility in public discourse over successive decades, pneumoconiosis persists as a predominant occupational malady on a global scale, with China being a notable epicenter, thereby inflicting considerable economic and public health encumbrances [3,4]. Such trends underscore the lacunae in extant policies intended for the amelioration of pneumoconiosis-related health risks. Consequently, there emerges an imperative to augment and refine current mitigation paradigms.

Through literature review on concurrent research, this essay will briefly introduce current medical diagnosis and treatments of pneumoconiosis, prevention of the disease, the global trend of pneumoconiosis and its contributing factors, policies aiming to mitigate health risks of pneumoconiosis from both government and private companies, delay between new policies and effectiveness, and propose an approach to shorten the delay.

2. General Information of Pneumoconiosis

2.1. Cause and Symptoms of Pneumoconiosis

In nations with a pronounced dependence on coal mining industries and coal as a primary energy substrate, the workforce within such sectors is disproportionately susceptible to heightened exposure risks [4]. The particulate-laden ambiance inherent to these industries substantially amplifies these hazards [4,5]. A preponderance of pneumoconiosis cases can be attributed to occupational exposures to specific aerosolized particulates, encompassing free silica, asbestos filaments, coal mine emanations, and heterogenous silicate particulates [4,5]. These aerosol contaminants are ubiquitously present in conventional sectors like coal extraction and thermoelectric power generation [3,4]. Intriguingly, in recent temporal epochs, incipient industries, namely nuclear fuel refinement and specialized equipment fabrication, have also manifested elevated risks associated with worker exposure to pneumoconiosis-inducing particulates [3]. Clinically, pneumoconiosis is delineated by protracted pulmonary inflammatory responses and subsequent lung fibrosis, both of which may culminate in profound physical incapacitation and mortality [1,5]. Pathognomonic features of this malady encompass sustained particulate exposure, chronic pulmonary inflammatory cascades, and ensuing pulmonary fibrotic transformations [1]. Silicosis stands as a quintessential exemplar of occupational pneumoconiosis, adversely affecting laborers in conventional domains such as mineral extraction, glass production, and in emergent sectors like synthetic stone fabrication and nanomaterial processing [3,5]. Recurrent exposure to crystalline silica particulates in occupational settings instigates chronic pulmonary inflammatory processes, which subsequently facilitate fibrotic lung alterations, ultimately manifesting as silicosis [5]. Clinical presentations of silicosis are typified by a persistent cough, dyspnea, and respiratory distress [5]. It is of paramount significance to underscore that pulmonary fibrosis, once initiated, is an irrevocable pathophysiological trajectory, rendering silicosis an exigent menace to both the pertinent industrial workforce and broader public health paradigms [1,5].

2.2. Diagnosis of Pneumoconiosis

The diagnostic process for silicosis and other forms of pneumoconiosis remains intricate and multifaceted [6]. Clinicians typically commence by eliciting detailed occupational histories, probing the duration and nature of particulate exposure such as coal dust and silica, as well as elucidating any symptomatic manifestations prior to delving deeper into diagnostic modalities [6]. Lung function tests are imperative to ascertain the functional integrity of the lungs, thereby facilitating an assessment of the disease's severity [7]. Notably, individuals afflicted with pneumoconiosis often exhibit hallmark lung function abnormalities, characterized by diminished total lung capacity, attenuated forced vital capacity, and compromised diffusion capacities [7]. In the wake of a strong clinical predilection for pneumoconiosis, radiographic examinations, specifically chest radiographs, might be requisitioned [6,7]. Yet, the diagnostic veracity of chest radiographs in discerning pneumoconiosis has been impugned in numerous scholarly deliberations [6]. For enhanced resolution and diagnostic precision, high-resolution computed tomography (HRCT) might be deemed necessary [6,7]. Albeit HRCT augments diagnostic accuracy, the financial implications, accessibility constraints, and potential radiation exposure temper its pervasive adoption [6]. Nevertheless, notwithstanding their inherent limitations, both chest radiographs and HRCT remain instrumental in the diagnostic algorithm for pneumoconiosis. These imaging modalities might unveil distinct pulmonary architectural aberrations, including but not limited to nodules, fibrotic patterns, or other parenchymal alterations indicative of inhaled particulate-induced injury [6]. For instance, the presence of diminutive, well-defined opacities could be evocative of simple pneumoconiosis, whereas more pronounced opacities or fibrotic changes might be redolent of progressive massive fibrosis (PMF), a more advanced and deleterious variant of the condition [6,7].

2.3. Treatments and Preventions of pneumoconiosis

In addressing the therapeutic strategies for pneumoconiosis, primary objectives gravitate towards symptom alleviation [8]. Foremost in the management paradigm of pneumoconiosis is the cessation of continued exposure to the inciting particulate matter [7]. This may necessitate occupational transitions or the instigation of enhanced safety protocols, including the deployment of protective facial coverings and the augmentation of workspace ventilation [2,7]. The pharmacological armamentarium for pneumoconiosis predominantly consists of bronchodilators and corticosteroids [8]. Bronchodilators are frequently administered to assuage bronchial muscular constriction, thereby facilitating unencumbered respiration, while corticosteroids aim to attenuate pulmonary inflammatory cascades [8]. As alluded to previously, the fibrotic sequelae in pneumoconiosis-afflicted individuals are irremediable [1,5]. Consequently, in the most grievous manifestations, pulmonary transplantation emerges as the ultimate therapeutic recourse [7].

Counteracting the pathophysiological onset of pneumoconiosis, preventive measures offer a more pragmatic avenue for mitigating its health ramifications. The etiology of pneumoconiosis is intimately tethered to protracted exposure to specific aerosolized particulates, encompassing free silica, asbestos filaments, coal mine emanations, and heterogenous silicate particulates [1,5]. Thus, precluding the inception of pneumoconiosis predominantly hinges on curtailing such exposures. From an individual protective standpoint, donning appropriate personal protective equipment, particularly facial masks or respirators, is pivotal [1,9]. It is imperative not only to select the apt respirator variant but also to ensure its meticulous fitment [1]. Concurrently, the edification of the workforce is paramount, fostering cognizance regarding the perils of particulate exposure, and underscoring the salience of protective gear and other prophylactic strategies [9]. On the infrastructural front, cultivating a workspace milieu that minimizes particulate dispersal and maintains optimal ventilation can be instrumental [9]. This might encompass the utilization of aqueous spray mechanisms, exhaust-driven ventilation, and other engineered interventions to curtail particulate liberation and airborne suspension. Periodic health surveillance and pulmonary function evaluations for those at heightened risk could facilitate the early detection of pneumoconiosis or its incipient markers, potentially obviating the dire ramifications of advanced disease progression [9]. In summation, it is incumbent upon employers to rigorously adhere to extant legal and safety statutes pertaining to workspace particulate exposures, fortifying the protective bulwark for their workforce.

3. Trend of Pneumoconiosis and its Contributing Factors

3.1. Global Trend of Pneumoconiosis and Re-emergence of Pneumoconiosis

While the global trajectory for newly diagnosed pneumoconiosis cases has been generally characterized by a decrement, a notable resurgence of the condition has been discerned in recent temporal epochs [2-4,10]. On a global scale, advancements in technology coupled with more stringent occupational health mandates have ushered in a declining prevalence of pneumoconiosis since the year 2015 [2,11]. The prevalence of coal worker’s pneumoconiosis has dropped from 23.33% before 1970 to around 2.29% in 2015[10]. Nonetheless, with the industrial ascension of emerging economies, there has been a concomitant proliferation in sectors such as mining, construction, and manufacturing, invariably augmenting the workforce's exposure to deleterious particulate matter [2]. In the annals of 2017, an alarming tally of over 60,000 new pneumoconiosis diagnoses were registered globally, culminating in an aggregate prevalence approximating 527,500 cases [11]. It is pivotal to emphasize that this revival of pneumoconiosis is not exclusively demarcated to developing countries. Within the context of developed countries, such as the United States, the frequency of pneumoconiosis, encompassing manifestations such as PMF, has been on an ascendant trajectory, with particular prominence in regions like central Appalachia as evidenced by data from 2019 [12]. Such epidemiological trends have galvanized governmental entities to recalibrate their strategies to attenuate the health ramifications of pneumoconiosis amongst the labor demographic and the broader

populace. These policy evolutions will be elucidated upon subsequently. However, one must remain cognizant of the potential temporal chasm that might ensue between policy promulgation and tangible policy efficacy.

3.2. Factors Contributing to Trend of Pneumoconiosis

A myriad of factors underpins the prevalence and resurgence of pneumoconiosis, with geographical characteristics of the workplace being a salient determinant [12]. In certain occupational settings, the act of rock excavation to access coal strata inadvertently exposes workers to rocks replete with silica concentrations, thereby elevating the risk of pneumoconiosis [12]. Illustratively, arid conditions might amplify particulate generation within sectors like coal mining, further exacerbating worker exposure [12].

Another dimension of this issue pertains to the potential temporal lag between the inception of policy and their outcomes, which will be talked later in detail. Such delays could inadvertently fuel the global recurrence of pneumoconiosis. Additionally, the advent of emergent industries delineates novel challenges for the mitigation of pneumoconiosis-associated health ramifications.

Notwithstanding the widespread acknowledgment of pneumoconiosis by governmental entities and entrenched industries, emergent sectors and the younger workforce remain inadequately apprised of potential pneumoconiosis hazards within their operational environments [3]. As previously emphasized, there is a palpable resurgence in pneumoconiosis episodes in recent years, spanning both developed and developing nations [2,10-12]. Another pivotal contributor is the rise of nascent industries, exemplified by sectors like nuclear fuel production and specialized equipment manufacturing [3]. In locales such as Tianjin, China, a cumulative 4,657 novel pneumoconiosis cases were chronicled from 2009 to 2018 [3]. Intriguingly, within this dataset, the mean duration of employment preceding the onset of pneumoconiosis symptoms in these emergent industries (19 years) starkly contrasts with their counterparts in traditional sectors (54 years) [3]. This discrepancy underscores either a pronounced deficit in pneumoconiosis cognizance or the existence of inherently high-risk environments within these burgeoning industries. Owing to this knowledge vacuum, particularly within emergent sectors, the workforce, predominantly the younger demographic, might be bereft of a comprehensive understanding of occupational health perils. Given the truncated occupational tenures observed in emergent sectors, it can be inferred that these personnel might be chronologically younger than their counterparts in conventional sectors. Consequently, these neophyte professionals in emergent arenas might possess a more limited repertoire of personal protective strategies, potentially owing to a dearth of experiential wisdom and mentorship from seasoned colleagues.

4. Response from Governments and Examples of Mitigation Strategies

4.1. Response from the US government

The resurgence of pneumoconiosis on a global panorama underscores an exigent need for the introspection and recalibration of extant workplace regulatory frameworks to more efficaciously mitigate associated health perils. In a seminal move towards this objective in 2016, the Occupational Safety and Health Administration (OSHA) embarked on a revision of the occupational exposure benchmarks pertaining to respirable crystalline silica (RCS), delineating a permissible exposure zenith of 50 mg/m3 and instituting more stringent stipulations for specific industrial domains [2]. This policy metamorphosis, which OSHA anticipates will substantially attenuate the likelihood of deleterious health outcomes, also integrated a comprehensive suite of safety tenets [2]. These encompass methodologies for the regulation of particulate concentrations, evaluative procedures for exposure quantification, deployment of tools engineered to curtail exposure, routine medical surveillance, and the meticulous archiving of pathological data [2,9]. As alluded to in prior discussions, a temporal hiatus might ensue prior to the discernible manifestation of the efficacy of newly minted regulations. Ergo, despite these commendable strides, the palpable renaissance of

pneumoconiosis on the global stage advocates for an exhaustive reexamination of the prevailing standards to ascertain their aptitude in addressing the concomitant health challenges. With greater specificity, novel benchmarks warrant institution in emergent industrial sectors, bolstered by rigorous investigative endeavors spotlighting potential aerial contaminants and concomitant health vulnerabilities intrinsic to such occupational milieus. Further, strategizing to truncate the latency between policy inception and its tangible impact emerges as a crucial vantage point.

4.2. Pneumoconiosis mitigation strategy – an example from the Chinese government

For optimal mitigation outcomes, collaborative endeavors from both governmental entities and the private sector are imperative. China, in its strategy to ameliorate the repercussions of pneumoconiosis, leverages its public insurance infrastructure to diminish the fiscal strain borne by afflicted workers [13]. Within the Chinese framework, the foundational medical insurance programs designated for urban employees, in conjunction with those tailored for rural inhabitants and non-employed urban denizens, encompass coverage for pneumoconiosis—especially in provinces evidencing pronounced incidence rates of the ailment. As articulated by the National Healthcare Security Administration of China, it was documented that, as of 2020, the foundational medical insurance apparatus would underwrite, on average, approximately 80% of the medical expenditures linked to pneumoconiosis [13].

4.3. Pneumoconiosis mitigation strategy – an example from the private sectors in China

Emerging from the private sector, the Huaibei Mining Holdings Corporation, a state-affiliated enterprise, has been assiduously furnishing its workforce with enhanced personal protective equipment and persistently disseminating knowledge pertaining to workplace safety [14]. Such endeavors have mitigated the ramifications of pneumoconiosis emanating from traditional industrial domains. Over time, the data gleaned from these initiatives holds potential to synergize academic and industrial pursuits, fostering the evolution of superior personal protective equipment and refining pedagogical guidelines on safety.

4.4. Pneumoconiosis mitigation strategy – the individual efforts

At an individual level, it is paramount for workers to remain apprised of the risks inherent to their occupational milieu and be cognizant of their entitlements [9]. With a nuanced understanding of such risks, laborers are poised to be more proactive in donning personal protective equipment and eschewing activities laden with potential hazards. Delving into their rights in an occupational context, familiarity with regional labor legislations and standards pertaining to occupational health can imbue workers with the agency to champion enhanced safety measures within their workplaces.

Yet, as previously delineated, the contemporary resurgence of pneumoconiosis underscores extant gaps in policy frameworks. This is particularly salient within the purview of nascent industrial sectors. Furthermore, the temporal chasm between the promulgation of regulatory edicts and their manifest efficacy remains a pressing conundrum.

5. Lag between Policy and Outcomes and One Proposed Approach

5.1. Temporal lag between inception of policy and outcomes

The translation of novel policy enactments into tangible outcomes within a pertinent domain is inherently characterized by temporal elongation before discernible effects materialize. This delay is shaped by a constellation of factors, each exerting differential impacts. Foremost, initiatives architected to modulate individual comportment may necessitate extended durations prior to witnessing widespread adoption among the target demographic. Drawing from the context of pneumoconiosis, although workers are inducted into the protocols of personal protective equipment utilization, there invariably exists a latency until this education achieves saturation, and workers

manifest a consistent proclivity to employ such equipment. Within this interstitial period, workers might already be subjected to pneumoconiosis-associated perils. Furthermore, even with instantaneous behavioral modifications, the consequent biological ramifications may not be immediate. Given the chronic nature of conditions like pneumoconiosis and the protracted exposure to etiological particulates, interventions devised to attenuate pneumoconiosis might only register a decline in morbidity metrics years, or potentially decades, subsequent, owing to the inherent latency of disease manifestation.

Additionally, certain policy directives are intertwined with infrastructural metamorphoses or capacity augmentation endeavors, which intrinsically demand protracted operational timelines. Illustratively, the establishment of optimally ventilated workspaces necessitates stages spanning design conceptualization, equipment procurement, and final installations, each introducing incremental temporal extensions. Measurement intricacies and data aggregation protocols further exacerbate this delay, as contingent upon the policy and the desired outcomes, pertinent data might only surface post a delineated period, thereby augmenting the perceptible latency.

Moreover, this delay is occasionally obfuscated by confounding elements. Exogenous events or concurrent policies might either amplify or attenuate the intended policy outcomes, challenging the unequivocal attribution of effects. Some policies might catalyze systemic perturbations, inducing feedback mechanisms that either accentuate or mitigate the envisioned outcomes. Consequently, the latency interceding policy inception and its discernible efficacy is a labyrinthine construct, exerting profound influences on the health trajectories of maladies like pneumoconiosis. Given the protracted average occupational tenures delineated both in traditional and emergent industries, as referenced in this exposition, any exacerbation of this delay potentially amplifies the burdens associated with pneumoconiosis. To this end, this essay proffers a potential stratagem to curtail this delay, specifically focusing on the facets of measurement and data acquisition.

5.2. Proposed approach – measurement and data collection

In emergent industrial sectors, the task of procuring precise data concerning atmospheric contaminants remains formidable until concomitant health repercussions surface conspicuously. Moreover, the inherent lag between policy promulgation and the realization of workplace safety protocols can potentially amplify health vulnerabilities within these sectors. Such challenges intrinsic to nascent industries might elucidate the observed renaissance of pneumoconiosis in contemporary times, signaling an urgent call for innovative mitigation paradigms. One such avant-garde proposition entails the conceptualization of a predictive model aimed at forecasting potential atmospheric contaminants derived from products within the workplace. Such a prognosticative approach could potentially attenuate the latency between policy inception and its tangible efficacy. By facilitating a generalized foretelling of potential workplace contaminants, the gap between regulatory edicts and their tangible translation into safety measures could be constricted, thereby mitigating adverse health outcomes. To buttress such regulatory endeavors, insights gleaned from data pertaining to exposure in traditional sectors could prove invaluable, lending credence to the feasibility of this approach.

Such endeavors necessitate a multidisciplinary collaboration spanning various societal strata. From a governmental perspective, the appropriate channeling of fiscal resources and the judicious allocation of pertinent departments are pivotal to undergird these innovative programs. Furthermore, efficacious dialogues between governmental bodies and private enterprises emerge as a cornerstone in this paradigm. Within the purview of the private sector, the furnishing of meticulous and accurate data from their operational environments is quintessential to the fruition of such initiatives, with their proactive participation in the rollout of new programs potentially yielding more immediate and pronounced dividends.

The academic sector also occupies a central role in this schema, particularly in terms of refining diagnostic techniques and devising novel interventions to mitigate pneumoconiosis's health implications. Innovations like biomarker assessments could truncate the aggregate medical expenditure associated with pneumoconiosis management. For instance, parameters such as the Mean

Corpuscular Hemoglobin Concentration (MCHC) exhibit a pronounced association with advanced stages of pneumoconiosis, positioning it as a potential diagnostic linchpin for future evaluations [15]. Furthermore, the academic community is entrusted with the onus of appraising health risks tethered to newly discerned atmospheric contaminants, providing indispensable advisement to both governmental and corporate stakeholders.

6. Conclusion:

In general, the global resurgence of pneumoconiosis underscores the urgent need for a comprehensive reevaluation of existing workplace regulation policies and the establishment of new standards in emerging industries. It is crucial to conduct thorough research to identify potential air pollutants and health risks in workplace environments of newly emerged industries. A combined effort from governments and private sectors is necessary to ensure better personal protective equipment, education on workplace safety, and adequate medical insurance coverage. On the personal ends, workers should also be aware about risks and personal protections. Additionally, innovative approaches, such as predicting potential ambient pollutants in the workplace, are required to shorten the delay between policy implementation and workplace safety measures, ultimately alleviating the impact of this debilitating disease.

References

[1] Li Y, Cheng Z, Fan H, Hao C, Yao W. Epigenetic Changes and Functions in Pneumoconiosis. Oxid Med Cell Longev. 2022;2022:2523066. Published 2022 Jan 20.

[2] Qi XM, Luo Y, Song MY, et al. Pneumoconiosis: current status and future prospects. Chin Med J (Engl). 2021;134(8):898-907. Published 2021 Apr 13.

[3] Zeng Q, Deng LH, Li XL, et al. Epidemiological characteristics of new pneumoconiosis in different industries in Tianjin. Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi. 2021;39(2):118-122.

[4] Wang D, Liang R, Yang M, et al. Incidence and disease burden of coal workers' pneumoconiosis worldwide, 1990-2019: evidence from the Global Burden of Disease Study 2019. Eur Respir J. 2021;58(5):2101669. Published 2021 Nov 25.

[5] Tan S, Chen S. The Mechanism and Effect of Autophagy, Apoptosis, and Pyroptosis on the Progression of Silicosis. Int J Mol Sci. 2021;22(15):8110. Published 2021 Jul 28.

[6] Li T, Yang X, Xu H, Liu H. Early Identification, Accurate Diagnosis, and Treatment of Silicosis. Can Respir J. 2022;2022:3769134. Published 2022 Apr 25.

[7] Krefft S, Wolff J, Rose C. Silicosis: An Update and Guide for Clinicians. Clin Chest Med. 2020;41(4):709-722. doi:10.1016/j.ccm.2020.08.012

[8] Adamcakova J, Mokra D. New Insights into Pathomechanisms and Treatment Possibilities for Lung Silicosis. Int J Mol Sci. 2021;22(8):4162. Published 2021 Apr 17.

[9] Jp NA, Imanaka M, Suganuma N. Japanese workplace health management in pneumoconiosis prevention. J Occup Health. 2017;59(2):91-103.

[10] Liu W, Liang R, Zhang R, et al. Prevalence of coal worker's pneumoconiosis: a systematic review and meta-analysis. Environ Sci Pollut Res Int. 2022;29(59):88690-88698.

[11] GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017 [published correction appears in Lancet. 2019 Jun 22;393(10190):e44]. Lancet. 2018;392(10159):1789-1858.

[12] Hall NB, Blackley DJ, Halldin CN, Laney AS. Current Review of Pneumoconiosis Among US Coal Miners. Curr Environ Health Rep. 2019;6(3):137-147. doi:10.1007/s40572-019-00237-5

[13] Reply from the National Medical Security Administration to Proposal No. 2998 of the Fourth Session of the 13th National People's Congress. Accessed September 29, 2023. http://www.nhsa.gov.cn/art/2021/10/12/art_110_7104.html.

[14] Huaibei Mining's efforts to prevent pneumoconiosis and care for workers. Accessed September 29, 2023. http://www.mohrss.gov.cn/gsbxs/GSBXSdifangxinxi/201704/t20170428_270141.html.

[15] Peng YF, Zhang QS, Luo WG. The Clinical Usefulness of Mean Corpuscular Hemoglobin Concentration in Patients with Pneumoconiosis. Int J Gen Med. 2023;16:3171-3177. Published 2023 Jul 27.