International Journal of Biology and Life Sciences
ISSN: 2957-9511 | Vol. 7, No. 3, 2024

Research Progress of COL5A1 in Osteosarcoma

Cheng Huang [1], Changshuai Qin [1], Zunian He [1], Xiaoban Huang [2], Changgong Lan [1, 2, *]

[1] Department of Orthopedic Surgery, The Affiliated Hospital of Youjiang Medical University for Nationalities, Baise, Guangxi, China
[2] Department of Infectious Diseases, The Affiliated Hospital of Youjiang Medical University for Nationalities, Baise, Guangxi, China

*** Corresponding author:** Changgong Lan (Email: landlong120@sina.com)

Abstract: COL5A1 (Collagen V alpha 1 chain) has gained significant attention as an important component of the extracellular matrix in recent osteosarcoma research. This review aims to summarize the expression characteristics, functional mechanisms, and potential clinical applications of COL5A1 in the context of osteosarcoma. By examining the latest findings, we delve into the role of COL5A1 in promoting osteosarcoma cell proliferation, migration, and invasion, while also assessing its potential as a biomarker and therapeutic target. Through this comprehensive analysis, we hope to provide valuable insights that can guide future research and clinical applications in this crucial area of oncology.

Keywords: COL5A1; Collagen V Alpha 1 Chain; Osteosarcoma; Extracellular Matrix; Biomarker.

1. Introduction

Collagen is an essential structural protein present in various connective tissues throughout the body, playing a key role in maintaining the integrity and functionality of these tissues. Its unique triple helix structure provides tensile strength and resilience, making it vital for the body's structural framework. Among the different types of collagen, collagen type V (COL5A1) is particularly important as it regulates cellular behaviors such as proliferation, migration, and differentiation, which are crucial for tissue repair and regeneration processes[1]. Beyond its structural role, collagen is also involved in signaling pathways that facilitate cellular communication and help maintain tissue homeostasis.[2].

Osteosarcoma is a malignant bone tumor that predominantly affects adolescents and young adults, posing significant clinical challenges due to its aggressive behavior and tendency to metastasize. The development of osteosarcoma is driven by a complex interaction of genetic, epigenetic, and environmental factors, which result in abnormal bone formation and tumor progression. Recent research has uncovered various molecular markers and signaling pathways that contribute to the development of osteosarcoma, highlighting the importance of high-mobility group AT-hook 2 (HMGA2) in facilitating tumor growth and metastasis[3]. Additionally, the tumor microenvironment is critical in the progression of osteosarcoma, with immune checkpoint molecules such as PD-1 and PD-L1 playing a significant role in modulating tumor immunity and influencing the effectiveness of therapeutic interventions[4].

The significance of COL5A1 in osteosarcoma research has attracted considerable attention due to its potential influence on tumor biology and its role as a prognostic marker. Research indicates that COL5A1 may affect the tumor microenvironment by altering extracellular matrix components and signaling pathways, which in turn can impact tumor behavior and patient outcomes[5]. This review aims to clarify the role of COL5A1 in osteosarcoma by examining its implications in tumor development and progression, as well as exploring potential therapeutic strategies. By delving into the molecular mechanisms that underlie COL5A1's involvement in osteosarcoma, we hope to provide valuable insights that could guide future research and clinical approaches to managing this complex malignancy.

2. Molecular Characteristics and Biological Functions of COL5A1

2.1. Structure and Gene Expression of COL5A1

COL5A1 is an essential part of the extracellular matrix (ECM) and plays a vital role in maintaining the structural integrity of various tissues. This gene, which encodes for COL5A1, is found on chromosome 9 and is expressed in multiple tissues, including skin, tendons, and ligaments. Its expression is carefully regulated during development and in response to mechanical stress, underscoring its significance in preserving tissue homeostasis. Research has indicated that changes in COL5A1 expression can lead to serious health issues, such as Ehlers-Danlos syndrome, where mutations in the COL5A1 gene result in a lack of collagen V, causing hypermobility and fragile tissues[6]. Additionally, COL5A1 has been linked to cancer progression, with increased levels associated with poor outcomes in various cancers, including breast and cervical cancers[7, 8]. The regulation of COL5A1 expression is influenced by intricate gene networks and signaling pathways, particularly those involving TGF-β, which is known to boost COL5A1 transcription in response to tissue injury and inflammation[9]. Gaining insight into the molecular mechanisms that control COL5A1 gene expression is essential for understanding its role in both healthy bodily functions and disease conditions.

2.2. Role of COL5A1 in the Extracellular Matrix

COL5A1 is crucial for the structure and function of the extracellular matrix (ECM), serving as a key component that supports tissue architecture. As a minor fibrillar collagen, COL5A1 plays an important role in the assembly of collagen fibrils, especially in combination with types I and III collagen, which are the main types found in connective tissues[10]. The presence of COL5A1 not only enhances the tensile strength and elasticity of tissues but also significantly impacts their mechanical performance. Furthermore, COL5A1 is involved in regulating various cellular behaviors such as migration,

proliferation, and differentiation by interacting with different ECM proteins and cell surface receptors[11]. Recent studies have shown that COL5A1 can also affect the tumor microenvironment, aiding cancer cell invasion and metastasis by altering the stiffness and composition of the ECM[12]. Additionally, its expression can change in pathological conditions like keloids and tendon injuries, where abnormal levels of COL5A1 can hinder healing and tissue remodeling[13]. Therefore, understanding the diverse roles of COL5A1 in the ECM is vital for developing therapeutic strategies that target ECM components in various diseases, including fibrotic disorders and cancer.

2.3. Expression Changes of COL5A1 in Osteosarcoma

2.3.1. Expression Level of COL5A1 in Osteosarcoma Tissue

COL5A1 is crucial for maintaining tissue integrity and influencing cellular behavior. In the context of osteosarcoma, a malignant bone tumor that primarily affects adolescents and young adults, research has indicated that the expression levels of COL5A1 are significantly altered when compared to normal bone tissues. Techniques such as quantitative PCR and immunohistochemistry have revealed that COL5A1 expression is notably upregulated in osteosarcoma tissues. This increase in expression may play a role in the tumor's aggressive characteristics and its capacity to metastasize. Additionally, the presence of COL5A1 within the tumor microenvironment can affect how cells interact with one another, creating a supportive niche that facilitates tumor progression. For example, COL5A1 has been linked to enhanced tumor cell proliferation and migration, suggesting its potential as a biomarker for diagnosing and predicting the prognosis of osteosarcoma. Moreover, the distinct expression patterns of COL5A1 in osteosarcoma compared to other sarcoma types imply that it may have a unique role in the pathophysiology of osteosarcoma[14, 15].

2.3.2. The Correlation between COL5A1 and Clinical Pathological Characteristics of Osteosarcoma

The clinical significance of COL5A1 in osteosarcoma goes beyond just its expression levels, as it has been linked to various clinicopathological features of the disease. Research shows that higher COL5A1 expression is associated with more advanced tumor stages, greater metastatic potential, and lower overall survival rates for patients. These findings imply that COL5A1 could act as an independent prognostic marker for osteosarcoma. Additionally, studies have examined the relationship between COL5A1 expression and the histological grade of tumors, finding that increased expression levels correspond to higher histological grades, which indicate more aggressive tumor behavior. Furthermore, the interaction between COL5A1 and signaling pathways involved in the progression of osteosarcoma, such as the TGF-β pathway, has been suggested, further linking COL5A1 to the disease's development. Gaining a deeper understanding of these correlations could help in creating targeted therapies that aim to modify COL5A1 expression or its activity within the tumor microenvironment, potentially enhancing treatment outcomes for patients with osteosarcoma[16, 17].

2.4. The Effect of COL5A1 on the Biological Behavior of Osteosarcoma Cells

COL5A1, a member of the collagen family, plays a crucial role in the behavior of osteosarcoma cells. Its expression levels are linked to tumor progression, metastasis, and overall patient survival outcomes. Increased levels of COL5A1 have been associated with various types of cancer, highlighting its potential as a biomarker for aggressive cancer characteristics. In the context of osteosarcoma, COL5A1 may influence the tumor microenvironment and affect how cells interact with one another, which in turn impacts the proliferation, migration, and invasion of osteosarcoma cells. Research indicates that higher expression of COL5A1 is correlated with poorer prognosis in multiple cancers, suggesting that it contributes to increased tumor aggressiveness and metastatic potential[18, 19].

2.4.1. The Effect of COL5A1 on Cell Proliferation

The influence of COL5A1 on cell proliferation in osteosarcoma has been demonstrated through various experimental approaches, showing that its overexpression can promote the proliferation of osteosarcoma cells. This effect may occur through the activation of signaling pathways that regulate cell cycle progression. For example, COL5A1 might interact with growth factors and components of the extracellular matrix, which can lead to increased cell survival and higher proliferation rates. Research has shown that elevated levels of COL5A1 correlate with enhanced tumor cell growth, potentially leading to worse outcomes for patients[11, 20]. Additionally, the expression of COL5A1 can be influenced by various non-coding RNAs, further underscoring its role in regulating cellular growth within the tumor microenvironment[12]. These findings highlight the importance of COL5A1 in osteosarcoma biology, suggesting that targeting this collagen could be a potential therapeutic strategy.

2.4.2. Regulation of COL5A1 on Cell Migration and Invasion Ability

COL5A1 plays a significant role in the migration and invasion abilities of osteosarcoma cells, which are essential for cancer metastasis. Research indicates that COL5A1 enhances cellular migration by improving the interaction between cancer cells and the extracellular matrix. This interaction aids in the breakdown of matrix components, enabling tumor cells to invade surrounding tissues [16]. Furthermore, increased levels of COL5A1 are associated with a rise in matrix metalloproteinases (MMPs), which are enzymes vital for remodeling the extracellular matrix and facilitating cancer cell invasion[21]. COL5A1 may also engage with various signaling pathways that control cell movement, thereby boosting the invasive capabilities of osteosarcoma cells. This invasive behavior not only leads to local tumor spread but also supports the formation of distant metastases, highlighting the potential of COL5A1 as a target for therapeutic strategies[12, 22]. Therefore, comprehending the function of COL5A1 in the migration and invasion of osteosarcoma cells is essential for developing effective approaches to tackle this aggressive form of cancer.

2.5. The Signaling Pathway and Mechanism of Action of COL5A1

2.5.1. Research Progress on COL5A1 Related Signaling Pathways

COL5A1 is an essential part of the extracellular matrix and has been linked to several signaling pathways that influence how cells behave, especially in cancer. Recent research has pointed out its involvement in activating the TGF-β signaling

pathway, which is crucial for controlling cell growth, differentiation, and programmed cell death. Variations in the COL5A1 gene have been connected to aortic dissection, likely due to the activation of this pathway, indicating its vital role in maintaining and remodeling blood vessels[23]. Moreover, COL5A1 appears to affect tumor development by enhancing communication between tumor cells and macrophages, which boosts inflammation and aids in the invasion of cancer cells[24]. Its role in response to low oxygen levels has also gained attention, as it serves as a prognostic marker in various cancers, suggesting it could be useful in clinical applications[19]. Additionally, COL5A1 interacts with different receptors in the ECM-receptor signaling pathway, significantly impacting the maintenance and differentiation of stem cells[25]. These insights highlight the critical role of COL5A1 in managing complex cellular responses and suggest its potential as a target for new therapies.

2.5.2. The Role of COL5A1 in the Tumor Microenvironment

The tumor microenvironment (TME) is a complex and ever-changing ecosystem that plays a crucial role in how cancer develops and responds to treatment. One of the key components within this environment is COL5A1, which significantly affects the mechanical characteristics of the extracellular matrix and influences the behavior of nearby cells. For example, in lung adenocarcinoma, COL5A1 produced by myofibroblasts has been linked to increased mechanical stress in tissues, which not only promotes the spread of cancer (metastasis) but also contributes to resistance against therapies[16]. Additionally, high levels of COL5A1 expression have been associated with poor outcomes in gliomas, suggesting it could serve as a valuable biomarker for tracking tumor progression[11]. The interaction between COL5A1 and immune cells in the TME is also important; it has been recognized as an immunological marker that can affect how the immune system responds to tumors[19]. Furthermore, COL5A1's involvement in the growth and spread of gastric cancer is highlighted by its regulatory interactions with lncRNA HOTAIR and miR-1277-5p, showcasing its diverse roles in cancer biology[26]. Overall, these findings about COL5A1's functions within the TME emphasize its significant impact on tumor behavior and highlight its potential as a target for new therapeutic strategies.

2.6. The Prospect of COL5A1 as a Biomarker and Therapeutic Target

2.6.1. Potential Application of COL5A1 in the Diagnosis of Osteosarcoma

COL5A1 has emerged as a significant biomarker in various malignancies, including osteosarcoma. Elevated levels of COL5A1 expression are linked to tumor progression and poor prognosis, indicating its potential utility in diagnosing bone sarcomas. Research shows that COL5A1 contributes not only to the structural integrity of the extracellular matrix but also to the dynamics of the tumor microenvironment, influencing cell behavior and interactions with immune cells. For example, studies have found a correlation between COL5A1 expression and the presence of tumor-infiltrating immune cells, which can impact the tumor's aggressiveness and its response to therapies[20]. Additionally, COL5A1's role in pathways related to cell adhesion and migration suggests its potential as a diagnostic marker for osteosarcoma, where early detection is crucial for improving patient outcomes. Its overexpression is associated with advanced disease stages, highlighting its relevance in clinical settings for stratifying patients based on risk and tailoring treatment approaches[11]. Therefore, COL5A1 represents a promising avenue for enhancing diagnostic accuracy and prognostic assessments in osteosarcoma, warranting further investigation into its mechanistic roles and clinical applications.

2.6.2. Research Progress and Challenges of COL5A1 Targeted Therapy

Targeting COL5A1 for therapeutic intervention offers a promising new approach in cancer treatment, especially considering its involvement in tumor progression and metastasis. Recent research has underscored the potential of COL5A1 as a therapeutic target, indicating that inhibiting it can hinder tumor growth and improve the effectiveness of current treatments. For instance, the RUNX1/miR-582-5p pathway has been recognized as a regulatory mechanism for COL5A1 in clear cell renal cell carcinoma, suggesting that targeting this pathway could disrupt the tumor-promoting effects of COL5A1[27]. Nevertheless, several challenges persist in the clinical application of therapies aimed at COL5A1. The diversity of tumor microenvironments and the intricate interactions of collagen within the extracellular matrix can complicate treatment outcomes. Furthermore, developing specific inhibitors that effectively target COL5A1 while sparing normal tissues poses a significant challenge[11]. Additionally, it is essential to comprehend the specific role of COL5A1 across different cancer types, as its expression may carry varying implications depending on the tumor context. As research advances, incorporating COL5A1-targeted strategies into clinical practice will necessitate thorough clinical trials to confirm their efficacy and safety, ultimately striving to enhance patient outcomes and treatment responses in cancers linked to COL5A1 dysregulation.

3. Conclusion

In conclusion, the role of COL5A1 in osteosarcoma is a vital area of research that highlights the complex relationship between genetic factors and tumor biology. Recent studies have shed light on the various ways COL5A1 contributes to the development of osteosarcoma, including its role in the composition of the extracellular matrix, its influence on cellular signaling pathways, and its interactions within the tumor microenvironment. Evidence indicates that COL5A1 serves not only as a structural element but also affects tumor cell behaviors such as proliferation, migration, and invasion.

As we continue to explore the evolving field of osteosarcoma research, it is important to reconcile the differing findings regarding COL5A1's function. Some studies emphasize its role in promoting tumors, while others suggest it may act as a tumor suppressor in certain contexts. This contrast highlights the need for a nuanced understanding of COL5A1 across different osteosarcoma subtypes and stages, underscoring the significance of context-dependent analysis in future research. Researchers must integrate findings from various models and methodologies to create a unified understanding of COL5A1's influence on osteosarcoma.

Looking forward, there is considerable potential for COL5A1 to be developed as a new clinical target for diagnosis and treatment. Given its involvement in tumor biology, therapies aimed at COL5A1 could improve existing treatment strategies and enhance patient outcomes.

Additionally, investigating COL5A1 as a biomarker for prognosis and therapeutic response could lead to more personalized approaches in managing osteosarcoma.

In summary, COL5A1 appears to play a significant role in the development and progression of osteosarcoma. Ongoing research is crucial to fully understand its intricate functions and to explore its potential applications in clinical practice. By promoting collaboration between researchers and clinicians, we can enhance our comprehension of osteosarcoma and refine treatment approaches, ultimately leading to better outcomes for patients impacted by this challenging disease.

References

[1] Lu X, Duan A, Liang S, Ma X, Deng T: Genomic Identification, Evolution, and Expression Analysis of Collagen Genes Family in Water Buffalo during Lactation. Genes (Basel) 2020, 11.

[2] Bronner-Shtrauchler O, Nativ-Roth E, Sanchez DS, Zaiden M, Vidavsky N: Multimodal characterization of the collagen hydrogel structure and properties in response to physiologically relevant pH fluctuations. Acta Biomater 2024, 178:170-180.

[3] Cui J, Dean D, Hornicek FJ, Yi G, Duan Z: Expression and Clinical Significance of High-Mobility Group AT-hook 2 (HMGA2) in Osteosarcoma. Orthop Surg 2022, 14:955-966.

[4] Hashimoto K, Nishimura S, Akagi M: Characterization of PD-1/PD-L1 Immune Checkpoint Expression in Osteosarcoma. Diagnostics (Basel) 2020, 10.

[5] Yang S, Liu L, Liu X, Li X, Zheng Y, Ren Z, Wang R, Wang Y, Li Q: The mitochondrial energy metabolism pathway-related signature predicts prognosis and indicates immune microenvironment infiltration in osteosarcoma. Medicine (Baltimore) 2023, 102:e36046.

[6] Colman M, Syx D, De Wandele I, Dhooge T, Symoens S, Malfait F: Clinical and molecular characteristics of 168 probands and 65 relatives with a clinical presentation of classical Ehlers-Danlos syndrome. Hum Mutat 2021, 42:1294-1306.

[7] Wu M, Sun Q, Mo CH, Pang JS, Hou JY, Pang LL, Lu HP, Dang YW, Fang SJ, Tang D, et al: Prospective molecular mechanism of COL5A1 in breast cancer based on a microarray, RNA sequencing and immunohistochemistry. Oncol Rep 2019, 42:151-175.

[8] Duan X, Ye D, Yuan J, Guan B, Guan W, Liu S, Fang J, Shi J, Zhu Y, Li Q, et al: COL5A1 overexpression correlates with poor prognosis in human cervical cancer. Int J Biol Markers 2024, 39:265-273.

[9] Sun M, Acosta AC, Emerick V, Adams S, Avila MY, Margo CE, Espana EM: Dysfunctional latent transforming growth factor β activation after corneal injury in a classical Ehlers-Danlos model. Matrix Biol 2024, 128:21-30.

[10] Machol K, Polak U, Weisz-Hubshman M, Song IW, Chen S, Jiang MM, Chen-Evenson Y, Weis MAE, Keene DR, Eyre DR, Lee BH: Molecular alterations due to Col5a1 haploinsufficiency in a mouse model of classic Ehlers-Danlos syndrome. Hum Mol Genet 2022, 31:1325-1335.

[11] Gu S, Peng Z, Wu Y, Wang Y, Lei D, Jiang X, Zhao H, Fu P: COL5A1 Serves as a Biomarker of Tumor Progression and Poor Prognosis and May Be a Potential Therapeutic Target in Gliomas. Front Oncol 2021, 11:752694.

[12] Dong C, Zhao Y, Yang S, Jiao X: LINC00173 blocks GATA6-mediated transcription of COL5A1 to affect malignant development of oral squamous cell carcinoma. J Oral Pathol Med 2023, 52:493-503.

[13] Lin P, Zhang G, Peng R, Zhao M, Li H: Increased expression of bone/cartilage-associated genes and core transcription factors in keloids by RNA sequencing. Exp Dermatol 2022, 31:1586-1596.

[14] Leiphart RJ, Weiss SN, DiStefano MS, Mavridis AA, Adams SA, Dyment NA, Soslowsky LJ: Collagen V deficiency during murine tendon healing results in distinct healing outcomes based on knockdown severity. J Biomech 2022, 144:111315.

[15] Leone MP, Morlino S, Nardella G, Pracella R, Giachino D, Celli L, Baldo D, Turolla L, Piccione M, Salzano E, et al: Specifications and validation of the ACMG/AMP criteria for clinical interpretation of sequence variants in collagen genes associated with joint hypermobility. Hum Genet 2023, 142:785-808.

[16] Zhu G, Wang Y, Wang Y, Huang H, Li B, Chen P, Chen C, Zhang H, Li Y, Liu H, Chen J: Myofibroblasts derived type V collagen promoting tissue mechanical stress and facilitating metastasis and therapy resistance of lung adenocarcinoma cells. Cell Death Dis 2024, 15:493.

[17] Jin L, Ma J, Chen Z, Wang F, Li Z, Shang Z, Dong J: Osteoarthritis related epigenetic variations in miRNA expression and DNA methylation. BMC Med Genomics 2023, 16:163.

[18] Feng G, Ma HM, Huang HB, Li YW, Zhang P, Huang JJ, Cheng L, Li GR: Overexpression of COL5A1 promotes tumor progression and metastasis and correlates with poor survival of patients with clear cell renal cell carcinoma. Cancer Manag Res 2019, 11:1263-1274.

[19] Zhu H, Hu X, Feng S, Jian Z, Xu X, Gu L, Xiong X: The Hypoxia-Related Gene COL5A1 Is a Prognostic and Immunological Biomarker for Multiple Human Tumors. Oxid Med Cell Longev 2022, 2022:6419695.

[20] Zhang J, Zhang J, Wang F, Xu X, Li X, Guan W, Men T, Xu G: Overexpressed COL5A1 is correlated with tumor progression, paclitaxel resistance, and tumor-infiltrating immune cells in ovarian cancer. J Cell Physiol 2021, 236:6907-6919.

[21] Wang C, Wang Y, Fu Z, Huang W, Yu Z, Wang J, Zheng K, Zhang S, Li S, Chen J: MiR-29b-3p Inhibits Migration and Invasion of Papillary Thyroid Carcinoma by Downregulating COL1A1 and COL5A1. Front Oncol 2022, 12:837581.

[22] Gong H, Wu W, Fang C, He D: CircBFAR correlates with poor prognosis and promotes laryngeal squamous cell cancer progression through miR-31-5p/COL5A1 axis. Laryngoscope Investig Otolaryngol 2022, 7:1951-1962.

[23] Chen P, Yu B, Li Z, Chen Y, Sun Y, Wang DW: COL5A1 Variants Cause Aortic Dissection by Activating TGF-β-Signaling Pathway. J Am Heart Assoc 2021, 10:e019276.

[24] Chen X, Ma C, Li Y, Liang Y, Chen T, Han D, Luo D, Zhang N, Zhao W, Wang L, Yang Q: COL5A1 promotes triple-negative breast cancer progression by activating tumor cell-macrophage crosstalk. Oncogene 2024, 43:1742-1756.

[25] Hu C, Zuo Q, Jin K, Zhao Z, Wu Y, Gao J, Wang C, Wang Y, Zhan W, Zhou J, et al: Retinoic acid promotes formation of chicken (Gallus gallus) spermatogonial stem cells by regulating the ECM-receptor interaction signaling pathway. Gene 2022, 820:146227.

[26] Wei Z, Chen L, Meng L, Han W, Huang L, Xu A: LncRNA HOTAIR promotes the growth and metastasis of gastric cancer by sponging miR-1277-5p and upregulating COL5A1. Gastric Cancer 2020, 23:1018-1032.

[27] Xue J, Zhu S, Qi F, Zhu K, Cao P, Yang J, Wang Z: RUNX1/miR-582-5p Pathway Regulates the Tumor Progression in Clear Cell Renal Cell Carcinoma by Targeting COL5A1. Front Oncol 2021, 11:610992.