

Ankle Lateral Collateral Ligament Injury Rehabilitation: Current Issues and Future Trends

Xiaoya Ma

Sports Training, Shenzhen University, Shenzhen, Guangdong, China

xyma925@gmail.com

Abstract. Lateral ankle ligament injuries are one of the most prevalent musculoskeletal injuries accounting for 40 percent of all sports-related trauma. Inadequate rehabilitation may result in a high incidence of re-injury and chronic ankle instability, which causes a lasting effects on mobility and quality of life. This paper discusses the anatomy of the lateral ankle ligament tears, existing rehabilitation evaluation techniques and management plans, and some of the obstacles and opportunities that define future clinical trends. The outcomes of the research show that the most vulnerable structure is ATFL and the primary subject of graded rehabilitation. Under the POLICE principle, functional rehabilitation has consistently outperformed as opposed to immobilization. Re-injury can be reduced through neuromuscular training and criterion-based return-to-sport protocols. High-grade injuries have yielded positive surgical results, particularly when there is particularly with early postoperative mobilization. The existing gaps are the inadequate prevention of chronic ankle instability, lack of individual protocols, psychological issues, a lack of specificity in exercises and long-term outcomes. Thus, a precision rehabilitation model comprising wearable technology, AI-assisted decision-making, and tailored individual care strategies holds strong potential to address current limitations and improve long-term patient outcomes.

Keywords: lateral ankle sprain, chronic ankle instability, rehabilitation, precision medicine.

1. Introduction

Lateral ankle ligament injury is one of the most prevalent musculoskeletal injuries worldwide, accounting for approximately 20 to 40 percent of all sports-related trauma. They are most frequent in activities involving jumping, sudden stops as well as rapid change of direction and are particularly common in sports like basketball, volleyball, and football [1]. The lateral collateral ligament complex is inherently vulnerable due to its anatomical structure: as the ankle is placed in plantarflexion and inversion position, the anterior talofibular ligament (ATFL) is placed under the greatest mechanical load, making it the most frequently ruptured structure and it is the most frequently ruptured structure [2].

In addition to the acute effect of pain and loss of function, lateral ankle ligament injuries have a serious long-term cost. Even in cases of well-supported populations recurrence is quite high, and a large proportion of the individuals develop chronic ankle instability (CAI), a condition characterized by a consistent feeling of giving way and recurrent re-injury that lasts over a year following the initial injury [3]. Reduced physical activities throughout the lifespan, impaired quality of life, and long-term potential of developing post-traumatic ankle osteoarthritis have been related to CAI [4]. This trend highlights the necessity of rehabilitation strategies that extend beyond control of the acute injury and actively work to prevent long-term disability.

Rehabilitation strategies across the globe have changed significantly in the last twenty years. Early functional rehabilitation as opposed to complete immobilization shows the current research momentum that endorses controlled loading as an effective technique in the healing of tissues and neuromuscular rehabilitation. A new principle of POLICE has replaced the RICE principle in acute management, and neuromuscular training of balance, peroneal reaction time, and proprioceptive recalibration has become a common feature of the rehabilitation continuum [3]. Imaging (especially MRI and dynamic ultrasound) has led to greater accuracy of characterizing injuries and allowed grading the severity more objectively [5]. The anatomical repair and its arthroscopic variants have

shown consistent return-to-sport outcomes in the surgical field, with growing evidence for increasingly mounting evidence of early postoperative mobilization protocols [6].

In China, there is a growing trend of incorporating traditional Chinese medicine (TCM) into practice in rehabilitation, such as acupuncture, tuina massage, and herbal compress therapy in addition to the Western physiotherapy. The integrative method is increasingly being supported by clinical evidence in the Chinese sports medicine literature, but high-quality randomized trial evidence for specific TCM components remains limited [7]. In a wider sense, the sports medicine rehabilitation system is yet to be established in the country, and the clinical studies published in this field are mostly rather short in their follow-up that restricts possibilities to gain understanding of long-term outcomes [7]. Specific literature on high-incidence sports like basketball and volleyball, which involve the jumping and cutting mechanics that are most closely linked with lateral ankle injury, is still limited.

To fill these gaps, this review examines the pathophysiology and biomechanical processes of lateral ankle ligament injuries by looking at their anatomy before analyzing the existing tools and techniques of rehabilitation assessment and treatment. The paper then identifies key contemporary problems in the rehabilitation practice and ends with a statement of the current technological and clinical trends that can change the future approach to care. The aim is to provide a comprehensive, evidence-based synthesis that can assist clinicians as well as researchers to move towards more individualized, effective, and sustainable rehabilitation.

2. Rehabilitation: Evidence, Challenges and Future Directions

2.1. Lateral Ankle Ligament Injuries: Anatomical Basis and Injury Mechanisms

2.1.1. Anatomical structures (ATFL, CFL and PTFL)

The lateral collateral ligament complex, which consists of the anterior talofibular ligament (ATFL), calcaneofibular ligament (CFL) and posterior talofibular ligament (PTFL) stabilizes the lateral ankle. The most frequently injured, the ATFL is between the anterior fibula and the talar neck and is biomechanically vulnerable in combined ankle plantarflexion and inversion. The CFL connects the distal fibula to the lateral calcaneus and passes through the tibiotalar and subtalar joints, hence playing a major role in stabilizing the two joints. Accordingly, a combination of ATFL and CFL tear leads to considerably more serious instability of the joint, compared to ATFL injury. The deepest and strongest structure, the PTFL is not easily affected in an ordinary sprain; it can only be traumatized in extreme situations such as severe ankle dislocations or rotational instability at the posterolateral position [1, 2].

2.1.2. Injury grading (Grade I to Grade III)

Lateral ankle sprains can be classified into three grades according to the levels of ligament disruption. Grade I trauma entails microscopic tears in the fibers in the absence of structural instability, mild swelling, and tenderness; recovery is typically achieved within one to two weeks with conservative treatment. Grade II injuries refer to partial macroscopic tears in the ligament, most commonly affecting the ATFL, which show evidence of increased swelling, ecchymosis, and laxity of the joint on stress testing. The duration of recovery takes between 2-6 weeks, and it is determined by the severity of the injury and quality of rehabilitation. Grade III injuries constitute complete rupture of one or more of the lateral collateral ligaments which present as evident joint instability and positive clinical symptoms (e.g., anterior drawer test). These acute injuries are likely to need surgery and extended rehabilitation [3, 4].

2.1.3. Biomechanical mechanisms

The ankle sprains are most prevalent in the case of the dynamic movements connected with the awkward landings, sudden stops, or swift direction changes, and physical collisions. The pathophysiologic process is focused on an excessive inversion moment produced by contact of the foot with the ground in a plantarflexed and inverted posture, and which is more than the load bearing

capacity of the lateral ligaments. Musculoskeletal simulation studies have validated that the ATFL experiences peak strain under jump landing and cutting maneuvers and hence it is the first to give way when placed under high-demand tasks. This is partly because the plantarflexed position preloads the ATFL, which further diminishes its failure threshold [8,9].

2.2. Rehabilitation Assessment System

The Ottawa Ankle Rules are used to initially assess acute ankle injury by imaging to identify the need to adopt radiography. In soft tissue assessment, MRI is the preferred tool and provides a detailed visualization of the tear in the ligament, damage of the cartilage of the articular cartilage and other soft tissue injury [5]. Additionally, ultrasound involves dynamic assessment in real-time when under stress, which makes it possible to observe and directly visualize the shift of the talofibular displacement. It has been demonstrated that the degree of talar displacement on ultrasound stress testing is a correlating factor in relation to injury severity and is a predictor of recovery time, making it a useful and cost-effective mode of initial assessment in an environment where MRI is not easily accessible [5].

2.3. Rehabilitation Interventions

2.3.1. Conservative treatment: physical therapy, neuromuscular training and bracing

The vast majority of Grade I and Grade II lateral ankle sprains are managed conservatively. Much of the old RICE (Rest, Ice, Compression, Elevation) principle had been replaced by POLICE (Protection, Optimal Loading, Ice, Compression, Elevation), and the paradigm shift is based on managed early weight-bearing and functional exercise, as opposed to the absolute immobilization. This paradigm shift is well supported by evidence: supervised functional rehabilitation interventions including progressive weight-bearing exercises with graduated exercises have been found to perform better than the passive methods in reducing pain, resolving the swelling, and returning to normal activity [3, 4].

The essence of the active rehabilitation is neuromuscular training, which involves progression starting with single-leg standing on the stable surfaces, moving on to unstable surface balance exercises and finally sport-specific movements (e.g., jumps, direction changes). The exercises address the fundamental deficiencies in the peroneal reaction time and the recalibration of the proprioceptive function. Semi-rigid ankle brace during the return-to-sport phase is highly advised since it offers further protection against re-injury, but it does not reduce the athletic performance significantly [1,7].

2.3.2. Surgical rehabilitation

Surgical management is recommended after conservative rehabilitation has failed to restore mechanical stability or when high-level athletes require a more predictable path back to competition. Anatomical repair in which a direct stitching of the torn ATFL and CFL with the addition of the inferior extensor retinaculum is done. Anatomical repair is preferred over non-anatomical reconstruction because it more closely preserves native joint biomechanics [10].

Postoperative rehabilitation has shifted toward earlier mobilization. In comparison with the patients confined to a cast over long periods after surgery, patients who start range-of-motion and weight-bearing exercises soon after surgery are demonstrated to experience better functional outcomes without an increase in complication rates. Return to the pre-injury sport level in professional athletes is possible in the majority of them after reconstruction and a standard duration is four to six months [9, 10].

2.3.3. Integrative Chinese-Western rehabilitation approaches

In the context of Chinese clinical practice, such a combination of conventional physiotherapy and traditional Chinese medicine (e.g., acupuncture, tuina massage, herbal compresses) can be integrated into the program of rehabilitation. These modalities focus on minimizing swelling in the area, reducing pain, and enhancing tissue healing, hence facilitating earlier engagement in active rehabilitation. Although there has been a growing support for integrated approaches in the Chinese

sports medicine community, the available high-quality randomized controlled trial evidence is still limited. To give further clarity on the best modalities of TCM, and when to implement it, and what clinical results are expected, further standardized studies are required [7].

2.4. Current issues

2.4.1. Inadequate prevention and management of chronic ankle instability

Chronic ankle instability (CAI), which affects approximately 20 to 40 percent of patients following a lateral ankle sprain, is characterized by persistent feelings of giving way and repeated re-injury beyond one year after the initial injury. In part, this can be explained by the fact that currently, common practice to clear athletes to return to activity after a specific period of time instead of on the basis of objective indication of functional recovery. Muscular abnormalities such as the retardation of peroneal response time and the inability to maintain posture are likely to remain after symptoms have disappeared, exposing the joint to re-injury [3, 11].

Evidence on professional football supports this with its high re-injury and time loss rates despite having access to dedicated medical resources. This suggests that the issue is not in the amount of resources but in the inconsistent application of criterion-based return-to-sport clearance. Clinicians should require athletes to meet objective standards of strength, balance and functional performance before they can resume full training [12].

2.4.2. Lack of individualized and precision rehabilitation protocols

One of the principal drawbacks of the existing practice is excessive dependence on one-size-fits-all and standardized rehabilitation protocols that lack consideration of differences in the severity of an injury, sport-specific demands of a patient, and individual biological responses. On the one hand, objective baseline assessment (e.g., imaging-acquired talar movement) has been demonstrated to influence predicting recovery time but is seldom taken into consideration to personalize rehabilitation programs during the first stage [6, 11]. Also, there is a wide range of clinical practice as some clinicians incorporate the concept of structured early functional rehabilitation, whereas others embrace passive methods which further contribute to diminishing the consistency and quality of care [11].

Ankle injury on the psychological level most notably due to kinesiophobia (fear of re-injury) is not normally taken seriously, yet may be a strong impediment to recovery. The fear may cause avoidance of movement, which deprives patients of the functional training needed to recover neuromuscular competence, and, consequently, slow recovery and increased chronic instability. Kinesiophobia and readiness to resume sport are among the psychosocial factors that need to be explicitly measured with the help of such tools as the Tampa Scale for Kinesiophobia (TSK) when it comes to rehabilitation testing. Strategies such as gradual exposure to challenging movements and structured goal-setting have demonstrated effectiveness in enhancing both confidence and functional outcomes [1].

2.4.3. Gap between research evidence and clinical practice

Many of the high-quality studies conducted on the rehabilitation of the ankle sprain process are mostly western based in Europe and North America, and may not directly have the same applicability to other health care systems, such as the developing sports medicine infrastructure in China. The majority of clinical research in China is of short-term nature, and as such, it cannot be extended to examine long-term effects like CAI development or post-traumatic arthritis. Moreover, sport-specific protocols are lacking for high-incidence sports such as (basketball, volleyball, football) where specific jumping and cutting mechanisms are engaged. General procedures can hardly serve these particular needs [7, 11] in an optimal way.

2.5. Future Perspectives

2.5.1. Rehabilitation decision-making based on artificial intelligence and big data

Applied to large clinical datasets, artificial intelligence has the potential to reveal trends that could guide more specific rehabilitation planning. Machine learning models might eventually identify which interventions produce better responses in specific patient profiles by processing the data of numerous patients such as their imaging results, assessment scores, psychological profiles as well as treatment results. This practice may practically be a clinical decision-support platform which produces customized rehabilitation plans depending on individualized patient baseline data, as opposed to the use of generalized protocols and clinician practice [7].

Biological augmentation strategies are a new option to the conventional treatment of patients that have severe or recurring ligamentous injury and who have not adequately responded to conventional treatment. The platelet-rich plasma (PRP) therapy involves the concentration of growth factors in the patient's own blood and the transmission of the factors to the injury area to help in repairing the tissue. Mesenchymal stem cell treatment goes one step further in which cells can potentially become ligament-forming fibroblasts and can contribute directly to structure regeneration. Though there is hopeful news associated with the preclinical evidence, the translation to well-designed large-scale human clinical trials continues to be a developing area, and these methods cannot be taken as standard care yet [7].

2.5.2. Precision medicine and individualized rehabilitation pathways

The general future trend in ankle ligament rehabilitation is that of a precision medicine model in which each element of the recovery program is tailored to the individual patient. This would include the combination of information from several sources: imaging, wearable sensors as well as functional performance tests and psychological screening. The rehabilitation stages would be determined by the realization of specific attained goals that are objective and measurable and not by the time elapsed. To scale up the realization of this vision, it will need improvements in data infrastructure, increased global cooperation concerning standardized outcome measures, and cultural transformation in clinical care toward data-based individualized care [1, 7].

3. Conclusion

Lateral ankle ligament injury is a clinically critical issue and optimizing rehabilitation outcomes requires a thorough understanding of the anatomical, biomechanical, and neuromuscular dimensions of the injury. As demonstrated in this paper, the ATFL is the weakest structure of the lateral ligament complex and the main target of both conservative and surgical rehabilitation. Active rehabilitation based on the POLICE principle and progressive neuromuscular therapy, in comparison to immobilization, has proven superior in all measures such as pain relief, swelling, and time to activity. Criterion-based return-to-sport protocols prove to be more effective compared with time-based methods to lower the occurrence of re-injury as well as preventing the onset of CAI. In Grade III injuries and refractory instability, anatomical repair done with early postoperative mobilization has good results, with most patients able to return to their pre-injury sport level. These results point to the importance of transitioning to an evidence-based and individualized model of rehabilitation. With the combination of objective data from imaging and functional tests with psychological screening and sport-specific requirements, clinicians may be able to formulate programs that are more consistent with the specific recovery requirements of each patient as opposed to just adhering to general guidelines. There are various limitations to present practice and research that should be mentioned. The vast majority of the high-quality evidence is produced in the Western environment and might not be applicable to every clinical situation. Existing literature has relatively short follow-up times, and there is a gap in knowledge about the long-term course of the results of rehabilitation, especially CAI incidence and post-traumatic degeneration of the joints. There exists very limited sport-specific research on the high-incidence sports like basketball and volleyball. Moving forward, wearable sensor

technology, artificial intelligence clinical decision support, and regenerative medicine solutions have promising potential to address these gaps. The combination of these tools into a precision rehabilitation system, whereby recovery pathways are customized on an individual basis by the use of objective, data-driven milestones, is the most promising way forward. To actualize this vision will require a long-term investment in longitudinal studies, international cooperation on standard outcome measures, and a more extensive cultural change in the direction of individualized, criterion-based care in any clinical environment around the world.

References

- [1] Martin RL, Davenport TE, Fraser JJ, et al. Ankle Stability and Movement Coordination Impairments: Lateral Ankle Ligament Sprains Revision 2021. *J Orthop Sports Phys Ther.* 2021;51(4):CPG1-CPG80. doi:10.2519/jospt.2021.0302
- [2] Aicale R, Maffulli N. Rotational ankle instability: A current concept review. *J Orthop Surg (Hong Kong).* 2023;31(2):10225536231182347. doi:10.1177/10225536231182347
- [3] Bergman R, Shuman VL. Acute Ankle Sprain. In: StatPearls [Internet]. StatPearls Publishing; 2026. PMID:29083595
- [4] Flore Z, Hambly K, De Coninck K, Welsch G. Time-loss and recurrence of lateral ligament ankle sprains in male elite football. *Scand J Med Sci Sports.* 2022;32(12):1690-1709. doi:10.1111/sms.14217
- [5] De Boer H, Robinson P. Imaging Findings in Acute Ankle Sprain and Syndesmosis Injury. *Semin Musculoskelet Radiol.* 2025;29(3):366-376. doi:10.1055/s-0045-1805077
- [6] Al-Janabi MM, Apostolides M, Southgate C, Dhinsa BS. Early mobilization following elective ankle lateral collateral ligament reconstruction in adults. *Foot (Edinb).* 2023;55:101988. doi:10.1016/j.foot.2023.101988
- [7] Yang Y, Wu Y, Zhu W. Recent advances in the management of chronic ankle instability. *Chin J Traumatol.* 2025;28(1):35-42. doi:10.1016/j.cjtee.2024.07.011
- [8] Shi X, Ganderton C, Adams R, Han J, El-Ansary D, Tirosh O. The past, present and future for chronic ankle instability research: A bibliometric analysis from 1992 to 2023. *Sci Sports.* 2025;40:491-502. doi:10.1016/j.scispo.2025.07.010
- [9] Boey H, van Rossom S, Verfaillie S, Sloten JV, Jonkers I. Maximal lateral ligament strain and loading during functional activities. *Clin Biomech (Bristol).* 2022;94:105623. doi:10.1016/j.clinbiomech.2022.105623
- [10] Maduka GC, Jakusonoka R, Maduka DC, Yusuf N. Conservative Management of Acute Lateral Ligaments of the Ankle Injuries. *Cureus.* 2023;15(10):e47709. doi:10.7759/cureus.47709
- [11] Goru P, Talha S, Majeed H. Outcomes and Return to Sports Following Ankle Lateral Ligament Reconstruction in Professional Athletes. *Indian J Orthop.* 2021;56(2):208-215. doi:10.1007/s43465-021-00532-0
- [12] Popp D, Weber J, Kerschbaum M, et al. Early functional treatment or trivialization? Current treatment strategies in lateral ligament injuries of the ankle. *Eur J Sport Sci.* 2021;21(10):1469-1476. doi:10.1080/17461391.2020.1845813